

**Andrew J. Riff, MD, Hinsdale Orthopaedic Associates S.C.**

**Locations:** 550 W Ogden Avenue, Hinsdale, IL 60521; 1010 Executive Court, Suite 250, Westmont, IL 60559; 2940 Rollingridge Rd #102, Naperville, IL 60564

**Phone:** For appointments and calls after business hours (8-4:30 M-F) please call 630-323-6116 (central scheduling and call forwarding). For clinical concerns during business hours, please call Dr. Riff's office (630-920-2314).

**Post-Operative Instructions – Shoulder Surgery**

**Prescription Meds:**

1. Norco (hydrocodone/acetaminophen): This is used for pain control. Take 1-2 tablets every 4-6 hours as needed for pain. Pain levels are highest on days 1 thru 3 post op, so you will want to take this on a regular schedule. After that, take as needed. Do not drive, drink alcohol or take Tylenol while taking this medication.
2. Colace 100mg (over the counter): Take two 100mg tablets (200mg), twice daily to prevent constipation after surgery.
3. Aspirin 81mg (over the counter): Take one 81mg tablet daily for 3 weeks to minimize your risk of a blood clot.

\*\*A regional block is placed which numbs your shoulder during surgery – this will wear off within 8-12 hours, plan to take pain medication throughout the early evening to ease the transition.

**Wound Care:** Leave the big surgical bandage on and do not shower for 48 hours after surgery. It is normal for the knee to bleed and swell following surgery. If blood soaks onto or through the dressing, this is not significant cause for concern. This is common for the first 24-48 hours after surgery. You may simply reinforce the dressing with gauze or change to a new clean dry gauze. After 48 hours, remove bandages.

**Bathing:** It is ok to shower after removing the dressing 48 hours after surgery. Let water run over Steri-strips, do not scrub. Do not submerge the wound in bathtub, hot tub, or pool for 4 weeks post-op.

**Physical Therapy:** Start physical therapy as soon as you are able to get in with a therapist 3 times a week.

**Sling:** Your sling should be worn for comfort for the first 2 weeks. You do not have to wear the sling if you don't feel the need.

**Ice Therapy:** It is very important to keep ice on your shoulder during the initial post-operative period (first 5-7 days). This should begin immediately after surgery. Use an ice machine continuously or ice packs (if no machine used) for 20-30 minutes every 2 hours daily until your sutures are removed. Care should be taken to avoid frostbite while icing by making sure the ice is not directly touching the skin.

**Follow-up:** You will need to follow up in clinic with Dr. Riff in 10-14 days. Please call central scheduling to make an appointment. At this visit you will receive updated instructions on your weight bearing, brace use and physical therapy.

**When should you contact the office?**

- If you have a fever >101.4 (a low grade temp is expected after surgery), but let us know if it gets this high!
- You develop chills or sweats
- You have pus or excessive bloody drainage from the wound, or if you have pain or redness surrounding the incision sites.
- You have nausea and vomiting that will not resolve with Zofran.
- Shortness of breath or chest pain