

Andrew J. Riff, MD, Hinsdale Orthopaedic Associates S.C.

Locations: 550 W Ogden Avenue, Hinsdale, IL 60521; 1010 Executive Court, Suite 250, Westmont, IL 60559; 2940 Rollingridge Rd #102, Naperville, IL 60564

Phone: For appointments and calls after business hours (8-4:30 M-F) please call 630-323-6116 (central scheduling and call forwarding). For clinical concerns during business hours, please call Dr. Riff's office (630-920-2314).

Post-Operative Instructions – Quad/Patellar tendon repair

Prescription Meds:

1. Norco (hydrocodone/acetaminophen): This is used for pain control. Take 1-2 tablets every 4-6 hours as needed for pain. Pain levels are highest on days 1 thru 3 post op, so you will want to take this on a regular schedule. After that, take as needed. Do not drive, drink alcohol or take Tylenol while taking this medication.
2. Colace 100mg (over the counter): Take two 100mg tablets (200mg), twice daily to prevent constipation after surgery.
3. Aspirin 81mg (over the counter): Take one 81mg tablet daily for 3 weeks to minimize your risk of a blood clot.

**Pain medication is injected into the wound and knee joint during surgery – this will wear off within 8-12 hours, plan to take pain medication throughout the early evening to ease the transition.

Wound Care: Leave the big surgical bandage on and do not shower for 48 hours after surgery. It is normal for the knee to bleed and swell following surgery. If blood soaks onto or through the dressing, this is not significant cause for concern. This is common for the first 24-48 hours after surgery. You may simply reinforce the dressing with gauze or change to a new clean dry gauze. After 48 hours, remove bandages. You do not need to replace them unless there is drainage.

Bathing: It is ok to shower after removing the dressing 48 hours after surgery. Let water run over Steri-strips, do not scrub. Do not submerge the wound in bathtub, hot tub, or pool for 4 weeks post-op.

Physical Therapy: You will initiate outpatient physical therapy beginning 2 weeks after the procedure.

Activity: You may bear full weight on your surgical leg. You should walk with your brace locked out straight. When at rest you may unlock it from 0-30 degrees to perform gentle range of motion. You should sleep with the brace locked straight.

Ice Therapy: It is very important to keep ice on your knee during the initial post-operative period (first 5-7 days). This should begin immediately after surgery. When using ice packs/ice bags, ice for 20-30 minutes with an hour off in between intervals.

Follow-up: You will need to follow up in clinic with Dr. Riff in 10-14 days. Please call central scheduling to make an appointment. At this visit you will receive updated instructions on your weight bearing, brace use and physical therapy.

When should you contact the office?

- If you have a fever >101.4 (a low grade temp is expected after surgery), but let us know if it gets this high!
- You develop chills or sweats
- You have pus or excessive bloody drainage from the wound, or if you have pain or redness surrounding the incision sites.
- You have nausea and vomiting that will not resolve with Zofran.
- Shortness of breath or chest pain